

# دراسة وحدة الضغط البسيطة، التوقعات النظرية والنتائج العملية.

د. محمد أبوبكر نوير الزوى      أ.د. محمد سليم الهاشمي.

## الملخص:

استخدام الضغط الهيدروديناميكي في تشكيل المعادن تمت دراسته بواسطة العديد من البحوث في السنوات الأخيرة، هؤلاء البحوث درسوا ظاهرة الضغط الهيدروديناميكي في عمليات السحب باستخدام الوحدات الهندسية المركبة والمستوية والمستدقة الطرف اللاإحتكاكية.

في هذه الورقة تم إجراء دراسة عملية لسحب الأسلاك بواسطة وحدة مستدقة الطرف لاإحتكاكية، واستخدام بوليمر Borosiloxane كمائع ضاغط.

نتائج هذه الدراسة تعتبر ذات أهمية في تصميم وزيادة كفاءة وحدة الضغط المستدقة الطرف لغرض السحب الهيدروديناميكي اللاإحتكاكي للأسلاك والأنابيب.

# Prediction and Experimental Results

M.A NWIR and M.S.J HASHMI<sup>1</sup>

*Department of Mechanical Engineering, Faculty of  
Engineering, P.O.Box 61160- HOON - Libya*

## ABSTRACT

The Application of Hydro-dynamic Pressure in Various Metal Forming Process has been studied by a number of research in the recent years. Theses researchers have studied the hydro-dynamic phenomenon in the drawing process by using a complex geometry, stepped or trapered bore pressure units. In this paper an experimental study as well as mathematical model for a simple taper pressure unit has been developed. A non-Newtonian pressure medium has been assumed. The results of this study should be useful in optimizing the design of the pressure unit for plasto-hydrodynamic die-less drawing and coating of wires, tubes and wire robes.

## INTRODUCTION

It has been observed experimentally [1, 2] that the pulling action of a wire through a viscous fluid gives a rise to drag force and generates hydro-dynamic pressure, the magnitudes of which depends on the type of fluid and the geometrical configuration of the pressure unit. A number of studies have been carried out applying this phenomenon to plastically reduce the diameter of copper wire. And tubes by drawing through pressure units of combined geometry, stepped and simple tapered bore. The first attempt to employ hydro-dynamic action in metal forming process [3] showed the development of hydro-dynamic action in wire drawing. Subsequently, a number of studies have been carried out applying the hydro-dynamic action in coating of wire, tubes, strips and wire ropes by using polymer melt. The use of polymer melt as the coating material was suggested in reference [4]. Limited experimental work in wire drawing and

---

<sup>1</sup>Prof M.S.J Hashmi School of Mech. & Manf. Eng. Dublin city University, Dublin 9, Rep. IRELAND

polymer lubrication investigating the coating features of polymer is given in reference [5], Whereas an in-depth study of die-less wire drawing process is given in reference [6]. Analytical solution presented in reference [7] for plasto-hydrodynamic pressure in a complex geometry pressure unit assuming Newtonian solution, Whereas in reference [8-9], experimental study to the distribution of hydro-dynamic pressure in a combined units is presented. A non-Newtonian plasto-hydro-dynamic analysis for the pressure generated due to the drawing of a wire through a simple tapered pressure unit in which brosiloxane [10] is used as pressure medium. The effects of shear rate and pressure on viscosity of the polymer together with the limiting shear stress are included in the analysis. The present analysis predicts the pressure distribution in the unit stress in the wire.




Fig. 1 Geometry used in the analysis

## EXPERIMENTAL PROCEDURES AND EQUIPMENTS

Experimental work was carried out using a purpose built each which incorporates the pressure unit and permits a wire speed of up to 0.7 m/sec. The schematic diagram of the drawing machine is shown in figure (2). The pressure unit which consists of a conical orifice, the melt chamber and the hopper are rigidly attached to the drawing machine bench. Details of the pressure unit is shown in the schematic

measure the hydro-dynamic pressure and drawing load respectively. The polymer melt chamber is attached to the pressure unit and is heated by electrical heater bands. The polymer was fed into the hopper in sufficient time was allowed for the polymer to reach the pre-set temperature. The diameter of the copper wire used is 2 mm. Borosiloxane was used as pressure fluid.

Before switching " on " the rig, the wire is guided through the melt chamber into the unit and passed over the pulley and then connected to the bull-block. The system is then checked for satisfactory drawing process. The hopper and the melt chamber are filled with enough borosiloxane for a complete set of test.




Fig. 2 View of the drawing bench

Thermo-couples are attached at appropriate positions so that the temperature can be monitored, controlled thermostatically and maintained at the pre-set level. Pressure transducers are connected to the recording and display units, the load cell is connected to the charge amplifier and then the rig is switched " on " and left for half an hour to reach the steady state temperature level. During the drawing process, the drawing load is noted from the chart recorder, the drawing speed is monitored by a hand held digital tachometer. The diameter of the uncoated wire is measured by a micrometer before

starting the test. At the end of each test, the motor, the chart recorder and the recording and display units are switched back to their original positions, test number is recorded on the data sheet and all the parameters are recorded for subsequent collection and analysis.




Fig. 3 Simple tapered unit

## RESULTS AND DISCUSSION

In obtaining the experimental results, tests were carried out by changing the inlet gap  $h_1$  and keeping  $h_2$  constant at 0.05 mm. Borosiloxane melt was used as the pressure medium and the melt temperature was kept at  $110 \pm 5$  °C for all the tests. For each combination of  $h_1/h_2$  gap ratios. Tests were performed at number of drawing speed ranging between about 0.05 and about 0.7 m/s.

Theoretical results were calculated based on the standard values of the parameters obtained from the experimental studies as following : initial viscosity  $\mu_0=50$  N.sec/m<sup>2</sup>, Length of the orifice,  $L = 180$  mm, Inlet gap,  $h_1 = 0.5$  mm, Final gap,  $h_2 = 0.05$  mm, Original diameter of the wire,  $D_0 = 2$  mm. The strain properties of the wire materials used have been determined and shown in figure (4) for the copper wire used.





Fig. 4 True stress strain curve



5.a pressure distribution curves




Fig. 5.b Theoretical pressure distribution curves

The Pressure distribution within the simple tapered unit based on experimentally measured pressure at three different points along the unit [fig. (3)], have been obtained and plotted for an over all gap ratios of  $h_1/h_2 = (3, 5, 10)$  and drawing speed of 0.1 m/sec as

given in figure (5). It is evident from the gap ratio considered, that the peak Pressure changes in magnitude and the point of occurrence with the change of the gap ratio and the drawing speed. For example, in figure 5.(a) for a gap ratio of  $h_1/h_2 = (3)$ , a maximum pressure of 450 bar was obtained, whereas, for a gap ratio of  $h_1/h_2 = 5$ , a higher magnitude of maximum pressure of 720 bar is observed, further increase to the gap ratio  $h_1/h_2 = 10$  gives lower magnitudes of maximum pressure of 550 bar.

Higher magnitudes of pressure were predicated theoretically as in whereas, in figure 5.(b), which illustrates pressure distribution for the same gap ratios and drawing speed, a higher magnitudes experimentally in figure 5.(a), and since the copper wire take place and results in relatively higher magnitude of the exit gap of the unit which in turn reduce pressure. It is therefore evident that elastic-plastic deformation takes place. Measurements of the dawn wire shows a reduction in the wire area.

To establish further the effectiveness of the pressure unit, experimental results were obtained in terms of the maximum pressure generated within the pressure unit.

The effect of drawing speed and the pressure ratio variation on the maximum pressure is illustrated in figure (6). The experimental results obtained in terms of maximum pressure versus drawing speed and gap ratio is given in figure 6.(a), which shows that a higher magnitudes of maximum pressure could be obtained by increasing the drawing speed. The effect of the gap ratio which is also included, indicates that there is a certain gap ratio at which a higher magnitudes of maximum pressure could be obtained.

In order to study the manner in which the maximum pressure is affected the magnitude of a number of parameters was varied systematically. In figure 6.(b) the inlet gap  $h_1$  was varied keeping  $h_2$  constant of 0.05 mm in order to obtain different gap ratios, the resulting maximum pressure for all drawing speeds reaches it's peak and then it remains fairly constant.

Results were also obtained experimentally and predicated theoretically in terms of drawing load for a different gap ratios and drawing speed. In figure (7), it is shown that for  $h_1/h_2$  ratio of 3 the drawing force was generally greater for all drawing speeds and that as this gap ratio increased drawing force of reduced magnitude are

obtained, both figures 7.(a) and (b) suggests that a generally higher magnitude of drawing load is obtained with lower gap ratios.




Fig. 6.a Drawing speed versus maximum pressure




Fig. 6.b Theoretical Drawing speed versus maximum pressure




Fig. 7.a Drawing speed versus drawing force




Fig. 7.b Theoretical Drawing speed versus drawing load

## CONCLUSION

The theoretical predications and the experimental results given suggest that optimum combination of gap ratio exist for the greatest magnitude of maximum pressure for a given drawing speed.



## REFERENCES

1. Hashmi M.S.J and Symmons G.R., "A numerical solution for the plasto-hydrodynamic drawing of a rigid non-Linearly strain hardening continuum through a conical orifice", Proc. 2nd Int. Conf. On numerical methods for linear problems, Barcelona, April 1984.
2. Hashmi M.S.J, Symmons G.R and Parvinmher H., "A novel technique of wire drawing", J.Mech. Eng Sci. Inst .Mech. Engrs 24 (1955).
3. Christopherson D.G and Naylor H., "Promotion of fluid lubrication in wire drawing", Proc. Inst. Mech. Engrs 169, 643-653 (1995).
4. Thompson P.J and Symmons G.R., "A Plasto-hydrodynamics analysis of the lubrication and coating of wire using a polymer melt during drawing", Proc. Inst. Mech. Engrs 191, 13, (1977).
5. Symmons G.R, Hashmi M.S.J and Parvinmher H., "Plasto-hydrodynamic, Die-less wire drawing : Theoretical treatment and experimental results", Proc. Int. Conf. Developments in drawing of metals Soc., pp. 54-62.
6. Parvinmher H, Symmons G.R and Hashmi M.S.J, " A non-Newtonian Plasto-hydrodynamic analysis of die-less wire drawing process using a stepped bore unit", Int. J. Mech. Sci. Vol. 29, No 4. Pp 239-257, 1987.
7. Al-Natour I and Hashmi M.S.J., "Development of a complex geometry pressure unit for hydro-dynamic coating applications", IMC-6, Dublin City University, Ireland, August/September 1989.
8. Nwir M.A and Hashmi M.S.J, "Plasto-hydro-dynamic Pressure distribution in a complex geometry pressure unit : Experimental results using Borosiloxane as Pressure Medium", IMC-11, The Queen's University of Belfast, August/September, 1994.
9. Nwir M.A and Hashmi M.S.J., "Hydro-dynamic pressure distribution in tapered stepped parallel bore and complex geometry pressure units: experimental results", AMPT'95, Dublin City University, Ireland, August, 1995 .00
10. Hull J.B, Jones A.R, Heppel A.R.W, Fletcher A.J and Trengove S., "The effect of temperature rise on the rheology of carrier media used in abrasive flow machining", Proc. Rude Int. Conf. On advances in coatings and surface engineering for corrosion

and wear resistance, University of Northumbria, Newcastle,  
U.K, May 1992, Vol 2, pp. 235-244.

# EVOLUTION

A Nonlinear Irreversible Quantum Model

*E. M. Elsheikh*

AL-TAHADI UNIVERSITY

## Abstract

Motivated by the postulates :

- (a)- Biological systems are compatible with the laws of quantum mechanics.
- (B)- Biological systems are order generating systems.
- (C)- Evolution is an irreversible transition from stable state to another state which is more structurally stable.

We discuss a definite model in which the life state of an organism, for successive generation, is a Schrodinger type of system which is nonconservation, nonlinear and irreversible. Vitality, the state variable, is a certain function of biological order.

Biological order, or complexity, is the quantity of information generated by growing organism. Whereas vitality, which is a function of the total energy metabolized by the organism and its life expectancy, is the capacity of the organism to generate biological order. Vitality has the following properties:

- i) It increases before adulthood, has a maximum when the organism is adult, decreases afterwards and becomes zero when the organism dies.
- ii) It is a periodic function of time with period,  $A$ , which is the life span of the organism .

We also discuss evolution within this model. We find that the evolution of a unicellular organism, being a process through which the life state function undertakes negative damping, leads to the increase of total vitality. This means that the evolution of unicellular organism leads to the increase of at least one of the following :-

- [A] Life span or cell cycle time.
- [B] Biological order or, complexity.
- [C] Energy consumption or body size.

## 1- INTRODUCTION :

A growing concern that the basic premises of the Darwinian theory of evolution, i.e. , random mutational changes and natural selection, are not sufficient to account for the enormous complexity and organization of present day biological systems has been gaining momentum since the mid of this century (Bertalanffy , 1954 ; Waddington , 1968 ; Gould 1982 ).

In April 1966 at Wistar institute of Anatomy and Biology in Philadelphia, a debate regarding this problem, took place between a group of mathematicians and biologists. The mathematicians charged that if natural selection has to choose from the astronomically large number of the alternative systems by means of the mechanisms described in current evolution theory the chances of producing a creature like ourselves is virtually zero. Eden (1966) who was especially concerned about the elements of randomness, contends that "No currently existing language can tolerate random changes in the symbol sequence which express its sentences. Meaning is almost invariably destroyed. Any changes must be syntactically lawful ones".

Thus all attempts to simulate the evolutionary process on computers have not been successful and the mathematicians concluded that current evolutionary theory is inadequate. It has to supply the programmer with a correct set of rules for " genetic grammaticality " that has a deterministic explanation rather than owing the observed stability of biological systems to selection pressure acting on random variations.

Whyte (1965) suggested that in addition to Darwinian selection there should be an internal selection at molecular, chromosomal and cellular levels, in accordance with their compatibility with internal coordination of the system. Whyte also asserted as some embryologists held, ontogeny is theoretically primary to phylogeny; consequently the synthetic theory can not be regarded as definitive until it has been combined with a theory of ontogeny. Waddington (1968) tried to show that evolution does not depend on random search. He emphasized that what occurs randomly are the mutations on the genome level, however, the output of these changes on the phenotype is not random, i.e. , there are certain operators that map the space of genotypes into a “ fitness space” .

Probably, based on Eden's (1966) objection to random search and on his assertion that “ No currently existing language can tolerate random changes in the symbol sequences which express its sentences “, Dawkin (1986) Proposed what he called cumulative selection. He clarified the concept by drawing attention to the fact that given a sentence of 28 units the probability that a monkey types it right away(What he calls single-step selection) is negligible. However, if whenever a letter falls in its proper place is preserved so that the next random change acts on the remaining letters then the chance for writing the sentence in this manner, which Dawkins calls cumulative selection, is very much improved. It is obvious that the process of generating a meaningful sentence in Dawkins above mentioned example, is guided by English Language grammar. However since such a grammar or set of rules to guide the evolutionary process have not yet been discovered Dawkins made some reservations as to significance of his example.

Some authors (Bertalanffy, 1954; Eden, 1965) even consider the evolutionary concept of fitness as tautologous, i.e, it restates the fact that only properties of organism that survive to produce offspring do survive. In other words to the question which organisms leave more offspring? The answer will be it is those that leave more offspring. So bertalanffy concluded that if leaving more offspring is the sole measure of evolutionary progress then it is difficult to see why evolution has progressed from the level of rabbits or even from bacteria.

Furthermore, from Darwin onwards, evolutionists realized that if we arrange all our available fossils in chronological order, they do not form a smooth sequence of gradual change. Most evolutionists, following Darwin, have assumed that this is mainly because the fossils record is imperfect as we thought. Maybe the gaps are a true reflection of what really happened, rather than being an imperfect fossil record. They suggested that evolution, maybe, in some sense goes in sudden bursts, punctuating long periods of stasis. Now accepting this punctuationistic pattern of evolutionary change necessitates in turn explaining why evolution occurred in this particular manner.

A radically different view which tries to overcome some of the above mentioned inconsistencies of neo-Darwinism was proposed by McClendon (1980). He compared biological evolution with the chemical evolution of isotopes. From the comparison he concluded that the forces which drive biological evolution are intrinsic property of matter. In other words the evolution of novel, more complex organisms, from lower ones precedes adaptation and selection.

The question whether evolution is a random process or an intrinsic property of matter is related to a deeper level of reality which concerns the nature of life. The nature of life has been the subject of heated debate, full with emotions, for centuries. The controversy culminated in present times to what may be called reductionism anti-reductionism dichotomy. Reductionists claim that biological phenomena are explainable in terms of physico-chemical laws, in practice the law of quantum mechanics. Antireductionists oppose this position and maintain that biological phenomena are autonomous and are irreducible to physics. If we accept the reductionists claim it will be difficult to explain why biological systems behave so differently from inanimate systems if they are both governed by the same set of laws. Likewise anti-reductionists are facing the fact that in the vast literature of biophysics or biochemistry there is no shred of evidence that the laws of physics are violated which implies the dependence of biological phenomena on physico-chemical laws.

To overcome this dilemma in order to pave the way for tackling the problem of evolution we need a physical theory of general

biology. We believe that the basic assumptions for such a theory were given by Pattee (1968) when he proposed that :

- A- Both living and non-living forms of matter obey precisely the same physical laws .
- B- Living states of matter are distinguishable from non-living states of matter only by the potential for evolution.

Starting from these assumptions, limiting our considerations to the laws of quantum mechanics, there are two possibilities for understanding life phenomenon based on the limitations we impose on the domain of quantum theory :

- A'- If we assume that the potential domain of quantum theory coincides with the domain of present day linear reversible quantum mechanics then we are left with no other alternative, due to the second assumption, than rejecting the dependence of life on laws and consequently try to reveal its mystery in terms of constraints, e.g non intergrable or measurement control constraints (Pattee 1990).

Of course Pattee did not assume that the domain of existing physical theory is closed. However, Pattee (1965) envisaged the development of physical theory in a particular manner when he said “ Notice that assumption A does not imply that all aspects of physical theory have been formulated, but only that whatever theories we currently accept must apply equally to living as well as nonliving matter”. Thus according to Pattee's program the evolution of physical or quantum theory is a continuous homogeneous process and not a hierarchical one. Such a view excludes the possibility that the potential domain of quantum theory which may account for life phenomenon may contain the set of present day quantum mechanics as a subset, i.e., Special case. This possibility which Pattee excludes is the only savior for his assumptions from contradiction as he did refer to some physicists who feel that his assumptions are contradictory (Pattee 1968).

is wider than present day linear reversible quantum mechanics then there is room for a nonlinear irreversible quantum mechanics. This is particularly evident if we conceive evolution as an irreversible transition from structurally stable state to another which is more structurally stable.

Since the development of any scientific theory, in this case quantum theory, is an open question and we can not claim a priori that the potential domain of the theory coincides with its present day existing boundaries, we shall take the view that life phenomenon, or at least evolution, is a nonlinear irreversible quantum phenomenon which is not reducible to present day linear reversible quantum mechanics. The irreducibility to the wave function is imposed by the second assumption which necessitates the distinction between animate and inanimate matter. However the proposed life state function cannot be quantum mechanical, i.e., Schrodinger type of system, and at the same time irreducible to the wave function. Unless, somehow, it admits a limiting transition to the wave function. This amounts to saying that linear reversible quantum mechanics is a subset or aspecial case of nonlinear irreversible quantum mechanics, so that reversibility is derivable from irreversibility and not vice versa .

What is the impact of this view on the long lived controversy between reductionists (mechanists) and anti reductionists (vitalists) ? It seems that, according to this solution, both mechanists and vitalists are partly correct. The mechanists are correct as far as they conceive life as a quantum phenomenon but they are not correct in claiming life reduction to present day linear reversible quantum mechanics. On the other hand the vitalists are correct when they oppose reducing life phenomenon to present day quantum mechanics but they are not correct in seeking life laws that are independent of quantum mechanics. Thus bearing in mind Pattee's assumptions we shall introduce the following as basic postulates for our model :

- (a) Biological systems are compatible with the laws of quantum mechanics.
- (B) Biological systems are order generating systems .



- (c) Evolution is an irreversible transition from structurally stable state to another state which is more structurally stable.

We also introduce "vitality", a function of the total energy metabolized by the organism and life expectancy, as a function of biological information. Whereby biological information is regarded as a measure of biological order or complexity. We require the vitality function to have the following properties :

- i) It increase before adulthood, has a maximum when the organism is adult decreases afterwards and becomes zero when the organism dies.
- ii) It is a periodic function of time with period A, which is the life span of the organism.

Consequently our system of postulates and conception of the vitality function satisfy the requirement that the life state of the organism can be described by a life state function which is a Schrodinger type of system, structurally stable, irreversible, has vitality as a state variable and admits a limiting transition to the wave function. We have confined, at this preliminary stage, our model to the evolution of unicellular organisms. It is found that evolution, being a process through which the life state function undertakes negative damping, leads to the increase of total vitality. The increase of total vitality means the increase of at least one of the following :

- [A] Life span or cell cycle time.
- [B] biological order or complexity.
- [C] Energy consumption or body size.

It is notable that in previous work the author (Elshiekh, 1987) disclosed some of these ideas :

## Definitions:-

2.11: Biological order, or biological complexity, is the quantity of information,  $I(t)$ , generated by a growing organism at time  $t$ ; where  $t$  is measured from the moment of initial growth.

2.12: Vitality,  $v(t)$ ; it is the capacity of the growing organism to generate biological order.

Information biologists may not have yet been able to calculate the information,  $I(t)$ , assigned to a growing organism. However from common physiological observation, at least qualitatively, the growing organism generates increasingly more information,  $I(t)$ , for  $0 \leq t \leq \alpha$ ; where  $\alpha$  is the time when the organism is fully grown, i.e., adult. Afterwards biological information,  $I(t)$ , decrease for  $t > \alpha$ . So it seems plausible to assume that  $I(t)$  has the following general properties :

$$\begin{aligned} I'(0) &> 0 \\ I'(\alpha) &= 0 \\ I''(\alpha) &< 0 \end{aligned} \quad (1)$$

So that the growing organism has maximum amount of information at adulthood. Now we shall construct a function  $v(t)$ , so that the relation :

$$I(t) = X[v(t)] \quad (2)$$

is valid for the general properties of  $I(t)$ . I.e., relations (1).

Let  $E(t)$  be the energy growth function defined as the energy available for the organism to support its growth, where  $t$  is the time measured from the moment of initial growth. For each organism we take  $F(t)$  to be defined by :

$$F(t) = (A - t)^a = 1^a \quad (3)$$

Where  $A$  is the life span of the organism,  $a$  is a positive parameter that depends upon the species of the organism and  $l$  is the life expectancy of the organism. We shall call the function  $F(t)$  life factor. The vitality function  $v(t)$ , is then defined by :

$$v(t) = E(t) F(t) = E(t) (A-t)^a \quad (4)$$

then  $v(A) = 0$

From Medawar (1945) and Bertalanffy (1957) we can deduce that the energy growth function,  $E(t)$ , for some classes of organisms, is given by :

$$E(t) = be^{Rt} \quad t \leq \alpha \quad (5)$$

Where  $b$  and  $R$  are positive constants which depend on the species of the organism and  $\alpha$  is the time, or age of the organism, when it is fully grown, i.e., adult. For the period  $\alpha \leq t \leq A$  we make the simple extension that metabolic rate,  $\frac{dE}{dt}$  remains constant at its value when  $t = \alpha$   
then,

$$E(t) = be^{R\alpha} + (t - \alpha) bRe^{R\alpha} \\ = be^{R\alpha} \{ 1 + R(t - \alpha) \} \quad \alpha \leq t \leq A \quad (6)$$

$$v(t) = be^{Rt} (A - t)^a \quad t \leq \alpha \quad (7)$$

$$v'(t) = be^{Rt} (A - t)^{a-1} \{ R(A - t) - a \} \quad t \leq \alpha \quad (8)$$

$$v(t) = be^{R\alpha} \{ 1 + R(t - \alpha) \} (A - t)^a \quad \alpha \leq t \leq A \quad (9)$$

Then  $v(A) = 0$ . And  $v(t)$  has single maximum for  $0 \leq t \leq A$ .

To restrict our model further we shall assume that  $v(t)$  takes this maximum value at adulthood, i.e., when  $t = \alpha$  so that  $v'(\alpha) = 0$ . This assumption, determine the value of  $a$  :

$$\begin{aligned} a &= R (A - \alpha) \\ &= E'(\alpha) L(\alpha) / E(\alpha) \end{aligned} \quad (10)$$

Note that  $a > 0$  as stated earlier since  $R > 0$  and  $A - \alpha > 0$ .

Consequently, in this model, there exists a vitality function  $v(t)$  that satisfies the following condition :

It increase before adulthood, has a maximum at adulthood, decrease afterwards and becomes zero when the organism dies .

## 2.2 - Numerical Example:

Using dimensionless variables we can draw the vitality curve. Let,

$$Rt = x, \quad AR = c, \quad R\alpha = c_0 \quad (11)$$

and

$$v(t) = \frac{by}{R^a}, \quad a = c - c_0, \quad \text{from eqn (10)}$$

$$y(x) = e^x R^a \left(A - \frac{x}{R}\right)^a = e^x (AR - x)^a = e^x (c - x)^a \quad (12)$$

$$x \leq R\alpha$$

$$x \leq c_0$$

From (9) :

$$\frac{by(x)}{R^a} = be^{c_0} \left\{ 1 + R \left( \frac{x}{A} - a \right) \right\} \left[ A - \frac{x}{R} \right]^a$$

$$y(x) = R^a e^{c_0} (1+x-c_0) \frac{1}{R} (c-x)^a$$

$$= e^{c_0} (1-c_0+x)(c-x)^a \quad (13)$$

$$\alpha \leq \frac{x}{R} \leq A$$

$$c_0 \leq x \leq c$$

Hence

$$y(x) = \begin{cases} e^x (c-x)^{c-c_0} & x \leq c_0 \\ e^{c_0} (1-c_0+x) (c-x)^{c-c_0} & c_0 \leq x \leq c \end{cases}$$

$c > c_0$ , take  $c = 8$  and  $c_0 = 4$

x	0	1	2	3	4	5	6	7	8
y	4096	6482	9590	12562	13759	8845	2620	218.4	0

It is worth mentioning that fig (I) is based on a particular form for the energy growth function, equations (5) and (6). However, it is evident, no matter what energy growth function the organism has, the present model always determines a vitality curve for it.

Now to substantiate relation (2) the function  $X(v)$  is constrained so that :

$$X'[v(0)]v'(0) > 0$$

$$\text{for } X'[v(0)] > 0 \quad (14)$$

$$X'[v(\alpha)]v'(\alpha) = 0$$

$$X''(v)v'^2 + X'(v)v''|_{t=\alpha} < 0$$

$$\text{for } X'[v(\alpha)] > 0$$

to obtain these conditions we have used (14).  
 $v''(\alpha) < 0$ . Noting that  $v'(0) > 0$  we see that conditions (14) are consistent with function  $X(x)$  that increase for  $x > 0$  from a vanishing value at  $x = 0$ . We have thus established that biological information,  $I(t)$ , is a function of vitality, or inversely we can say that vitality is a function of biological information which is measure of biological order.

The following definitions will be useful in our subsequent considerations

Instantaneous total vitality  $V(t) = \int_0^t v(x) dx$  (15)

Total vitality  $V(A) = \int_0^A v(t) dt$  (16)

Average vitality density for all ages,

$$\bar{v} = \frac{\int_0^A v(t) dt}{A} = \frac{V(A)}{A} \quad (17)$$

Average Instantaneous total vitality,

$$\bar{V}(t) = \int_0^t v(x) dx = \bar{v}t \quad (18)$$

### 2.3 - Successive Generations:

The model may be usefully employed to discuss vitality for successive generations. We shall essentially be concerned with unicellular organisms particularly those which reproduce by binary fission. For such systems fig (I) represents the average vitality for one generation, i.e., when the organisms dies. However a unicellular organism usually does not die, it starts to divide when it is fully grown. Referring to fig (i) division occurs at time  $t = A_1$ , Where

$\alpha < A_1 < A$ . The parent cell gives birth to two identical daughter cells so that  $v(A_1) = 2v(0)$  where  $v(0)$  is the initial vitality. Again each daughter cell grows and divides in the same manner. Thus the average vitality function, vitality per unit cell or organism, or successive generations  $v_g$ , under constant environmental conditions is a periodic function of time with period  $A$  (assuming no confusion we have deleted the subscript from  $A_1$ ).

Then,

$$v_g = v(t + nA) = v(t). \quad (19)$$

Where  $n = 1, 2, 3, \dots$  is the number of cell divisions or generations. Figure(ii) represents this state of affairs :

Having been fully grown the unicellular organism divides.

If the unicellular organism completes the cell cycle and dies instead of mitoting then fig(i) describes such a situation, i.e., determines the vitality curve. However for fig(ii) to be possible the organism must divide successively, i.e., must have a natality rate  $r(t)$ . Accordingly we shall regard fig(ii) representative for the life state and that vitality,  $v$ , and natality,  $r$ , are the fundamental biological state variables.

Where,

$$r(t) = \frac{dp}{dt} / p \quad (20)$$

$p$  is the population size.

Of course the life state of a biological systems being natality or multicell-ular is affected by factors other than vitality & natality, c.g. environmental factors.

So, how do we justify not incorporating these as separate state variables? The answer is that, within the present model, we assume all effects on the life state caused by such factors to occur by virtue of their effects on vitality and natality only. In other words we assume that vitality and natality are functions of these factors, and the effect of these factors is to be interpreted as a perturbation or a modification of vitality and natality. This is the same argument by which Rosen

(1976) justified why in the kinetics of a chemical reaction the state variables to the rate constants and not incorporating factors such as temperature, pressure and volume as separate state variables.

### 3.1 - Life State Function:

In search for a plausible modular life state function we introduce the following postulates:

Postulate (b):

Biological systems are order generating systems.

Based on this postulate, i.e., on the fact that biological systems are order generating systems, and on the fact that vitality, being a function of biological order, is a state variable it is natural, referring to fig.(ii), to represent the life state of the unicellular organism by what we shall call life state function (L). Then,

$$L = L(v) \quad (21)$$

where  $v = v_a$

To complete the definition of L we introduce :

Postulate (a):

Biological systems are compatible with the laws of quantum mechanics.

This postulate derives its strength from the fact that there is no shred of evidence any-where in the vast literature of biochemistry or biophysics that the laws of quantum mechanics are invalid (Elsasser, 1981) and from Von Neumann's mathematical result that there can be no second set of laws independent of quantum mechanics.

Now from postulate (a) and equation(21) it is reasonable to assume that the life state function, L, is a Schrodinger type of system with vitality as a state variable. We shall therefore take for it the simple form :

$$L = L(0)e^{i\omega/h} \quad (22)$$



where  $L(0)$  is the amplitude,  $h$  is planck's constant. Now according to eqn. (21),  $\Phi$  must be a function of  $v$ . Further we must have correct dimensionality. With this in mind we limit our considerations to the concrete example in which  $\Phi$  is given by:

$$\Phi = \int_0^t E(1 - \frac{x}{A})^a dx \quad (23)$$

this completes the definition of  $L$ . From equations (4), (15) and (23) we obtain :

$$\Phi = A^{-a} V(t) = A^{-a} \int_0^t v(x) dx \quad (24)$$

$$\begin{aligned} \therefore L &= L(0) \exp. \frac{1}{A^a h} \int_0^t v(x) dx \\ &= L(0) \exp. \frac{1}{A^a h} \int_0^t E(A-x)^a dx \end{aligned} \quad (25)$$

The life state function, equation (25), has important and desirable features, namely, the differential expression of equation (25) is

$$L'' - \frac{v'}{v} L' + \frac{v^2}{k^2} L = 0 \quad (26)$$

where  $k = h A^a$

Using (21) we can also have:

$$L'' - \frac{(L')^2}{vL'(v)} + \frac{v^2}{k^2} L = 0 \quad (27)$$

Equation (27) is that of a non-conservative non-linear system which maintains its stability by consuming energy from the surroundings. Consequently the oscillations given by fig. (ii), Which

oscillations and that the system is structurally stable. A result which is compatible with empirical observation. Moreover, it is significant that, using the vitality function, we can determine the life state function without undertaking the impossible task of constructing the Hamiltonian for the whole organism. By so doing we shall gain a new insight and account within our model for some aspects of biological phenomena. Finally, we would also like to emphasize the significance of the periodicity of the life state function by referring to the enormous efforts that have been made to reveal the oscillatory nature of fundamental biological organization in general and cellular performance in particular, (Goodwin, 1963; Nicolis, 1977; and Yates, 1981).

The Irreversibility of the life function is also evident since the vitality function equation (4) can not accept any substitution of  $-t$  for  $t$ .

Now using (17):

$$\bar{v} = \text{constant } t \quad (28)$$

$$\bar{v}' = 0 \quad (29)$$

we shall reduce (26) to the simpler form :-

$$L'' + \frac{\bar{v}^2}{k^2} L = 0 \quad (30)$$

we note that:

$$\text{Frequency, } w = \frac{\bar{v}}{k} \quad (31)$$

$$\text{Wavelength, } \lambda = \bar{v}A = V(A) \quad (32)$$

where, from equation (16),  $V(A)$  is the total vitality of the system.

#### 4.1 - Evolution:

The life function (25) is constructed to describe the life state under constant environmental conditions. However, we know that environmental conditions do not always remain constant, they change. In doing so the organism interacts and sometimes incorporates these changes and perturbations. The most significant perturbations from an evolutionary point of view are mutational changes. These perturbations induce nonlinearity in the system. Accordingly the system may spiral to a focus due to positive damping in which case the mutation is lethal and the system may extinct. On the other hand the system may exhibit stable periodic solution in the neighborhood of the homogenous solution with greater period due to negative damping in which case the mutation is beneficial. Thus we say that the system has evolved. The terms positive and negative damping are used in electrical engineering in connection with oscillatory phenomena which dissipate and absorb energy respectively. Likewise we use these terms to characterize the dissipation and generation of vitality by the system when it interacts with the environment. Thus, to incorporate evolution in our model, we make the following assumption:

Evolution ( mutation, selection, ... etc) of unicellular organism is a process through which the life state function undertakes negative damping.

#### Result:

The evolution of the unicellular organism leads to the increase of its total vitality,  $V(A)$ .

#### Proof :

Let the unicellular organism before evolution be given by:

$$L'' + \omega^2_1 L = 0 \quad (33)$$

has undertaken negative damping. The new evolutionized system becomes :

$$L'' + \omega_2^2 L = \varepsilon f(L, L') \quad (34)$$

Where  $\varepsilon$  is a small positive parameter that characterizes the smallness of the deviation of  $\omega_2^2 L$  from  $\omega_1^2 L$ . If the deviation is not small we set  $\varepsilon = 1$ . Biologically the fact that  $\varepsilon \neq 0$  accounts for certain generation of vitality which did not exist when  $\varepsilon = 0$ . Being interested in finding a stable periodic solution in the neighborhood of the homogenous solution, under the condition of negative damping, equation (34) is readily solvable by Krylov and Bogoliubov (1947) asymptotic method for damping oscillation. The solution yields :

$$\omega_2 < \omega_1 \quad (35)$$

Since, in general, the wavelength,

$$V(A) \propto \frac{1}{\omega}$$

then,

$$V_2(A) \propto \frac{1}{\omega_1}, \quad V_1(A) \propto \frac{1}{\omega_2} \quad (36)$$

Where  $V_1(A)$ ,  $V_2(A)$  are the total vitality of the organism before and after evolution respectively. Equations (35) and (36) yields,

$$V_2(A) > V_1(A) \quad (37)$$

Which proves our result.

Equation (37) proves that the evolution of the unicellular organism leads to the increase of its total vitality. However we need to know what does the increase of total vitality mean ?.

From equation (37), (16) and (14) we obtain :

$$\int_0^{A_2} (E_2 L_2^a) dt > \int_0^{A_1} (E_1 L_1^a) dt \quad (38)$$

From (38), also referring to fig (I), the increase of total vitality leads to the increase of at least one of the following :-

[A] Life span or cell cycle time, A.

[B] Biological order or complexity,  $v(\alpha)$ .

[C] Energy consumption or body size,  $E(\alpha)$ .

Looking at the phylogenetic evolution of uni-cellular organisms, e.g. , evolution of bacteria, we encounter features such as increase of life span and body size which are in conformity with the present model.

### 5.1 - Limiting Transition:

The irreducibility of the life function, begin a Schrodinger type of system, necessitates a limiting transition to linear reversible quantum mechanics. Accordingly given the life function we obtain:

$$\frac{-ih}{L} \frac{dL}{dt} = \frac{E(t) (A - t)^a}{A^a} \quad (39)$$

This looks like Schodinger's equation with a time dependent energy. It therefore appears plausible to characterize inanimate matter with time independent energy on the right-hand-side of this equation.

In order words we need to add " inanimate energy"  $\epsilon$ , the energy that the body of the dead organism will have as a lump of matter.

Then (39) will read when modified :

$$\frac{-ih}{L} \frac{dL}{dt} = \frac{v(t)}{A^a} + \epsilon \quad (40)$$

for living organisms, so that :

$$\frac{L}{dt} = \epsilon \quad (41)$$

for inanimate matter, in the latter case one may identify  $L = \psi$  so that

$$-i\hbar \frac{d\psi}{dt} = \epsilon\psi \quad , \text{i.e.,} \quad H\psi = \epsilon\psi \quad (42)$$

Where  $H$  is the hamiltonian and (42) is Schrodinger's equation.

## 5.2 - Summary :

We have, within a concrete model, discussed various concepts that relate to the growth, reproduction, vitality, information, biological order, structural stability, irreducibility and evolution of unicellular organisms. We see that our system of postulates and conception of the vitality function necessitate the description of biological evolutionary process by a life state function, a proposed order generating principle, which is nonlinear, nonconservative and irreversible Schrodinger type of system. Thus random mutational changes generate beneficial mutations because the system is capable to increase its total vitality through negative damping, otherwise the probability of generating beneficial mutations is virtually zero.

The logical structure of our model is based on the view that the total domain of quantum theory is a hierarchical set and not a homogeneous one, so that quantum laws which govern biological phenomena admit limiting transition to linear reversible quantum mechanics. In other words the quantum biological laws contain the physics of inanimate systems, i.e., what is called ordinary laws of physics, as a limiting case. Such a view, of course, has a tremendous impact on the question of reduction. It seems it is no longer significant or relevant to ask whether biology is reducible to physics or vice versa. Because we have reached the limit where the domain of theoretical biology coincides with the domain of theoretical physics. This is simply because we have got one Universe.

It is worth mentioning that the model is potentially capable to account for problems related to ontogeny. e.g. , we can envisage cellular differentiation as a process through which the life state function undertakes negative damping; because the whole organism is usually issued from a single cell, zygote, through successive cellular divisions. Accordingly ontogeny recapitulates phylogeny in the sense that cellular differentiation is also a process of increase of total vitality, i.e., increase of life span or cell cycle time and / or body size.

Finally the present model can easily be generalized to account for multicellular organisms.

#### Acknowledgments:

I would like to acknowledge my indebtedness to Professor M.O Taha for his substantial contribution to this work. I am also grateful to Professor B.C. Goodwin for helpful comments.

## REFERENCES

1. Bertalanffy, L. Von, 1954. "Problems of Life". Academic Press New York
2. Bertalanffy, L. Von, 1957. "Quantitative Laws in Metabolism and Growth". *Quart. Rev. Biol.*, 32:217-231.
3. Dawkins, R. 1986. *The Blind Watchmaker*. Longman Scientific and Technical Group. U.K.
4. Eden, M. 1966. "Inadequacies of New Darwinian Evolution as a Scientific Theory", In (Moorhead, S.P. and Kaplan, M.M eds.). *Mathematical Challenges to the New-Darwinian Interpretation of evolution*. The Wister Institute Press.
5. Eldredge, N., and Gould, J.S.. 1972. *Punctuated Equilibria: An Alternative to Phyletic Gradualism*. In T.J.M Schopf, et., *Models in Paleobiology*, 85-115. San Francisco: Freeman, Cooper.
6. Elsasser, M.W.1981. "Principles of a New Biological Theory A Summary. *J.Theor. Biol.* 89, 131-150.
7. Elsheikh, M. 1987 "Evolution of Unicellular Organisms in Terms of Vitality Function." *Proceeding of the Third International Congress of Biomathematics*. Santiago, Chile.
8. Goodwin, C.B.1963. "Temporal Organization in Cells: A Dynamic Theory of Cellular Control Process". Academic Press London & New York.
9. Gould, S.J., 1982. "DARWINISM and the Expansion of Evolutionary Theory". *Science* 21 : 380-387.
10. Krylov, N.N. Bogoliubov, 1947. "Introduction to Non-linear Mechanics." Princeton N.J : Princeton University Press.



11. McClendon J.H. 1980 .” The Evolution of the Chemical Isotopes as and Analog of Biological Evolution.” *J.Theor. Biol.* 87, 113-28.
12. Medawar, P.B. 1945, “Essays on Growth and Form”. Gross Clark and Medawar P.B. (eds).
13. Nicolis, G., and Prigogine, I-1977. “Self Organization in Non Equilibrium Systems: Form Dissipative Structures to Order Through Fluctuations. Wiley-Interscience Publication John Wiley & Sons. New York, London.
14. Pattee, H.H 1968, The Physical Basis of Coding & Reliability in Biological Evolution. In Waddington, ed., Towards a Theoretical Biology. The Kynock Press. Birmingham.
15. Rosen, R 1970. “Dynamical System Theory in Biology “. Vol Wiley Interscience, NewYork, London.
16. Waddington, C.H. 1968. “Does Evolution Depend on Random Search?” In (Waddington, ed.) Towards a Theoretical Biology. The Kynock Press Birmingham.
17. Whyte, L.L. 1965. “ Internal Factors in Evolution” Associated Book Publisher Limited.
18. Yates, E.F. 1982. “ Outline of a Physical Theory of Physiological Systems”. *Can. J. Physio. Pharmacal.* Vol-60.

